

## Section 1.5

### Exponential Functions

General Form of an exponential function  $f(x) = ca^n$

Where  $a^n = \underbrace{a \cdot a \cdot a \cdot a \cdot a \cdot a \cdot \dots \cdot a}_{n\text{-times}}$

Example  $6^4 = 6 \cdot 6 \cdot 6 \cdot 6 = 1296$

Laws of Exponents

1)  $a^x a^y = a^{x+y}$

2)  $\frac{a^x}{a^y} = a^{x-y}$

3)  $(a^x)^y = a^{xy}$

4)  $(ab)^x = a^x b^x$

5)  $a^{\frac{m}{n}} = (\sqrt[n]{a})^m = \sqrt[n]{a^m}$

6)  $a^0 = 1$

---

### Example 1

a) Simplify  $(2x^4)(5x^8)$

Solution:  $(2x^4)(5x^8) = 10x^{4+8} = 10x^{12}$

b) Simplify  $(5x^3)^2$

Solution:  $(5x^3)^2 = 5^2(x^3)^2 = 25x^6$

c) Simplify  $16^{\frac{3}{2}}$


Solution:  $16^{\frac{3}{2}} = (\sqrt{16})^3 = 4^3 = 64$

---

---

## Graphs of exponential functions

Exponent curves are sometimes referred to having a “hockey stick shape”  
Here are some examples of exponential functions.




---

### Example 2

Graph the following function  $f(x) = 2^x$


To the graph the function we will use the strategy of choosing five x values and making a table of values.

$x$	$f(x)$
-2	$f(-2) = (2)^{-2} = \frac{1}{2^2} = \frac{1}{4}$
-1	$f(-1) = (2)^{-1} = \frac{1}{2^1} = \frac{1}{2}$
0	$f(0) = (2)^0 = 1$
1	$f(1) = (2)^1 = 2$
2	$f(2) = 2^2 = 4$



**Example:** Graph the following function  $f(x) = 2^{-x}$

$x$	$f(x)$
-2	$f(-2) = 2^{-(-2)} = 2^2 = 4$
-1	$f(-1) = (2)^{-(-1)} = 2^1 = 2$
0	$f(0) = (2)^0 = 1$
1	$f(1) = (2)^{-1} = \frac{1}{2^1} = \frac{1}{2}$
2	$f(2) = (2)^{-2} = \frac{1}{2^2} = \frac{1}{4}$



---

## The Exponential Function

The Euler Number:  $e \approx 2.718$

---

### Example 3

a) Find  $e^2$

Solution:  $e^2 = 7.39$

b) Find  $e^{-3}$

Solution:  $e^{-3} = .050$


---

## Graphs of the exponential function

### Example 4

Graph  $f(x) = e^x$

$x$	$f(x)$
-2	$f(-2) = e^{-2} = \frac{1}{e^2} = .14$
-1	$f(-1) = e^{-1} = \frac{1}{e^1} = .37$
0	$f(0) = e^0 = 1$
1	$f(1) = e^1 = 2.7$
2	$f(2) = e^2 = 7.4$





---

### Example 5

Graph  $f(x) = e^{-x}$


$x$	$f(x)$
-2	$f(-2) = e^{-(-2)} = e^2 = 7.4$
-1	$f(-1) = e^{-(-1)} = e^1 = 2.7$
0	$f(0) = e^0 = 1$
1	$f(1) = e^{-1} = \frac{1}{e^1} = .37$
2	$f(2) = e^{-2} = \frac{1}{e^2} = .14$



---

### Example 5


Graph  $f(x) = 3^x - 2$




---

### Exponential Modeling

#### Exponential Growth



## Exponential Decay



---

### Example 6

If a bacteria population starts with 200 bacteria and double every three hours, then the number of bacteria after  $t$  hours is  $B(t) = 200(2)^{\frac{t}{3}}$

a) Find the population of the bacteria after 15 hours?

$$B(15) = 200(2)^{\frac{15}{3}} = 200(2^5) = 200(32) = 6400$$

b) Find the population of the bacteria after 25 hours?

$$B(25) = 200(2)^{\frac{25}{3}} = 200(322.54) \approx 64508$$